



## How to assess gas concentrations in laying hens farms?



### Legislation

#### Directive 98/58 EC, Annex, Paragraph 10:

“[...] gas concentrations must be kept within limits which are not harmful to the animals.”



### Method




#### Description of the method:

The method involves measurements of concentrations of ammonia (NH<sub>3</sub>) and carbon dioxide (CO<sub>2</sub>) using a gas meter.



#### Preparation for each inspection:

- Make sure that the gas meter has been calibrated within the last six months in a certified laboratory.
- Make sure that the gas meter measures 0 ppm NH<sub>3</sub> outdoor in open air.



©ANSES



#### During the inspection:

Number of points measured in the barn:

**≥8,000 birds:** six measures in total, with two measures per one-third of the barn.

- In multitier systems: three points on the floor, two points at the first tier and one point at the second tier.
- In single-tier systems: three points on the floor, three points on the slats/tier.

**<8,000 birds:** four measures in total, with two measures in each half of the barn.

- In multitier systems: two points on the floor, two points at the first tier.
- In single-tier systems: two points on the floor, two points on the slats/tier.

For each measure, the gas meter is held still at the height of the birds' heads for 60 s, after which the gas concentration is noted.

When choosing the specific measuring point:

- Take the airflow from the ventilation into account.
- Make sure that some measures are near feed and water lines.
- Avoid the wettest areas.

When measuring CO<sub>2</sub> concentration:

- Make sure not to exhale in the direction of the gas meter.
- Do not measure directly under the heating sources.

# How to assess gas concentrations in laying hens farms?

## After inspection:




- Disinfect the gas meter.
- Let the gas meter run in open air outdoor until it shows 0 ppm NH<sub>3</sub> before being used during a new on-farm inspection. This eliminates all NH<sub>3</sub> accumulated in the sensor during the previous inspection.


## Interpreting the data

From interviews of official veterinarian inspectors from different Member States, it appears that there are two main ways of using the measurements:

- 1) Calculate the average of the six measures and use the mean to decide whether the gas concentration complies with the legislation.
- 2) Use the highest measure to decide whether the gas concentration complies with the legislation.

## Uncertainties/reservations

-  Gas concentrations fluctuate within a day, between days and between seasons. The method described above is a point observation valid only for the time of measuring.
-  The gas meter is costly and needs regularly calibration for a sufficiently high level of reliability.
-  There are differences between models and brands of gas meters in accuracy, so before purchasing gas meters this should be examined. The accuracy has been shown to be between 1% and 15% when the measurement is carried out on a reference gas (a mixture of CO<sub>2</sub> or NH<sub>3</sub> in pure air)<sup>1</sup>.



©ANIS

## Sources

A number of sources have been searched for protocols on how to measure gas concentration in poultry barns. These include scientific peer-reviewed articles and documents collected from competent authorities in Member States within EU and from non-EU countries within Europe. The methods considered to have the highest validity and reliability, while still being feasible during on-farm inspections, were found in protocols from the French<sup>5</sup> and Swiss<sup>6</sup> competent authorities. The present method has been developed based on these protocols.

<sup>1</sup> Précision de la mesure des concentrations en gaz. The French Poultry Technical Institute (ITAVI) and Institut National de la Recherche Agronomique (INRA).

<sup>2</sup> Protocol 'DGAL/SDSPA/2017-998', Direction générale de l'alimentation, Service des actions sanitaires en production primaire, Sous-direction de la santé et de protection animales, Bureau de la protection animale (BPA), France.

<sup>3</sup> Protocol 'Luftqualität', Division of Animal Welfare, Federal Food Safety and Veterinary Office FSVO, Switzerland.